

Necropsy: Respiratory Disease Sampling Guidelines

The ideal samples for investigation of respiratory diseases are those that are collected within 4-8 hours of death, however, any sample is better than none, and the lab does not charge based on the number of samples submitted. **Both** fresh and formalin fixed tissues from **2-4** different sections of lung including the junction between normal and diseased tissue as well as the following samples are recommended for submission:

Organ	Formalin*	Fresh	Size Fresh Samples
Heart ⁺	Yes	Yes	4 x 4 x 2 cm
Lung ⁺ (2-4 sections)	Yes	Yes	4 x 4 x 2 cm
Trachea ⁺	Yes	Yes	4 x 4 x 2 cm
Bronchial Lymph Node +	Yes	Yes	4 x 4 x 2 cm
Mesenteric Lymph Node	Yes	Yes	
lleum	Yes	Yes	10 - 20 cm length

^{*} A 2 x 2 x 1 cm piece of tissue from the organs listed above should be placed in formalin. The ratio should be 9 parts formalin to 1 part tissue for proper fixation. The WVDL recommends using a 16 ounce wide mouthed jar that contains 8 ounces of 10% buffered formalin (Fischer catalog # 23-426-797). Since the formalin jars tend to leak; **please place the formalin jar inside a Ziploc® bag**.

Shipping Requirements

- Veterinarians should completely fill out the Necropsy Submission Form which includes the
 age of the animal. An electronic copy is available at www.wvdl.wisc.edu. Click on the
 Submission and Order Forms link to download the submission form.
- The laboratory should receive the samples within 24-36 hours so samples collected on Friday should be refrigerated and shipped on Monday.
- Package the samples with a sufficient number of ice packs to ensure they remain cold during shipment.
- Check out our special shipping rates here.
- **WINTER SHIPMENTS:** Please add isopropyl alcohol (1:10 ratio) to the formalin for formalin fixed tissues to prevent freezing during shipment.

CL-RES-37 Active Date: 12/1/24

⁺ Denotes essential tissue for submission.